


AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions and listings of claims in the application:

1. (Currently amended) A method of treatment **for Type II** for diabetes and its complications and associated conditions, comprising administering compounds **selected from** of Formula (1) (Gibberellins)



Formula (1)

wherein

R¹ is H or a group -O-R²⁰, where R²⁰ is H, a glycosylic ether group (glycoside ether), C₁₋₆ alkyl group, or R¹ together with R² or R¹⁰ forms a bond (C₁-C₂ or C₁-C₁₀ double bond, respectively);

R² is H or a group -O-R²¹, where R²¹ is H, a glycosylic ether group (glycoside ether), or together with R⁴ forms a bond (lactone) or R² together with R¹ or R³ forms a bond (C₁-C₂ or C₂-C₃ double bond, respectively);

R^3 is H, =O, or $-O-R^{22}$, where R^{22} is H or a glycosylic ether group (glycoside ether), or R^3 together with R^2 forms a bond (C_2-C_3 double bond);

R^4 is OH, or $-OR^{23}$, where R^{23} is unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, arylalkyl, amidine, $-NR^{24}R^{25}$ or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur; R^{24} and R^{25} may or may not be the same, are hydrogen, or C_{1-20} alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulphur; or R^4 together with R^{21} or R^{28} forms a bond (lactone);

R^5 is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted C_{1-20} alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters;

R^6 is H or OH or together with R^7 forms a bond ($C_{11}-C_{12}$ double bond);

R^7 is H, =O, or $-OR^{26}$, where R^{26} is H or a glycosylic ether group (glycoside ether) or R^7 together with R^6 forms a bond ($C_{11}-C_{12}$ double bond);

R^8 is H, hydroxyl, mercaptan, or halogen, amino, azido, $NR^{24}R^{25}$, unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, or arylalkyl, or $-OR^{27}$, where R^{27} is a glycosylic ether group (glycoside ether);

R^9 is H or OH, or together with R^{15} forms a bond (C_9-C_{15} bond);

R^{10} is H, CH_3 , CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH_2O-R^{28} or $-OR^{28}$, where R^{28} is H or together with R^4 forms a bond (lactone) or R^{10} together with R^1 forms a bond (C_1-C_{10} double bond);

R^{11} is H, or OH or is absent;

R^{12} is CH_3 , CH_2OH , $COOH$ or a glycosylic ester (glycoside ester) of said $COOH$;

R^{13} is methylene, or a divalent hetero-atom, or NR^{29} , where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl; and a double bond is present between C_{16} and R^{13} when R^{11} is absent; or R^{13} is H, OH, CH_3 , CHO, CH_2X , where X is halogen, $CHNR^{29}$ where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl when R^{11} is H or OH; with the proviso that where R^{11} is OH, R^{13} is not OH;

R^{14} is H or OH;

R^{15} is H, or together with R^9 forms a bond (C_9 - C_{15} bond);

and its and/or pharmaceutically acceptable **lactones, esters, active esters, salts and organic bases,** derivatives to a patient in need thereof.

2. (Original) The method of claim 1, wherein the complications and associated conditions of diabetes are one or more of: obesity, micro and macro vascular diseases, nephropathy, neuropathy, eye diseases, and diabetic ulcerations.

3. (Original) The method of claim 1, wherein the Gibberellins are Gibberellin A_3 .


4. (Original) The method of claim 1, wherein the Gibberellins are a mixture of Gibberellin A_3 and Gibberellin A_4 and/or Gibberellin A_7 .

5. (Currently amended) The method of claim 1, wherein the pharmaceutically acceptable ~~derivatives are salts~~ **are selected from** including alkali metal salts, alkaline earth metal salts, metal, and salts of ammonium **[[,]]** or organic bases.

6. (Original) The method of claim 5, wherein the organic bases are lidocaine, or $\text{NR}^{16} \text{R}^{17} \text{R}^{18} \text{R}^{19}$, where R^{16} , R^{17} , R^{18} , R^{19} , which may be the same or not the same, are hydrogen, or substituted or unsubstituted C_{1-20} alkyl, alkanol, or aryl groups.

7. (Canceled)

8. (Currently amended) A method of treatment **for Type II** of diabetes and **its complications and associated** related conditions comprising administering an ~~effective amount of a compound~~ **selected from** of formula (1) (Gibberellins)



Formula (1)

wherein

R^1 is H or a group $-\text{O}-\text{R}^{20}$, where R^{20} is H, a glycosylic ether group (glycoside ether), C_{1-6} alkyl group, or R^1 together with R^2 or R^{10} forms a bond (C_1-C_2 or C_1-C_{10} double bond, respectively);

R^2 is H or a group $-\text{O}-\text{R}^{21}$, where R^{21} is H, a glycosylic ether group (glycoside ether), or together with R^4 forms a bond (lactone) or R^2 together with R^1 or R^3 forms a bond (C_1-C_2 or C_2-C_3 double bond, respectively);

R^3 is H, =O, or $-O-R^{22}$, where R^{22} is H or a glycosylic ether group (glycoside ether), or R^3 together with R^2 forms a bond (C_2-C_3 double bond);

R^4 is OH, or $-OR^{23}$, where R^{23} is unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, arylalkyl, amidine, $-NR^{24}R^{25}$ or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur; R^{24} and R^{25} may or may not be the same, are hydrogen, or C_{1-20} alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulphur; or R^4 together with R^{21} or R^{28} forms a bond (lactone);

R^5 is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted C_{1-20} alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters;

R^6 is H or OH or together with R^7 forms a bond ($C_{11}-C_{12}$ double bond);

R^7 is H, =O, or $-OR^{26}$, where R^{26} is H or a glycosylic ether group (glycoside ether) or R^7 together with R^6 forms a bond ($C_{11}-C_{12}$ double bond);

R^8 is H, hydroxyl, mercaptan, or halogen, amino, azido, $NR^{24}R^{25}$, unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, or arylalkyl, or $-OR^{27}$, where R^{27} is a glycosylic ether group (glycoside ether);

R^9 is H or OH, or together with R^{15} forms a bond (C_9-C_{15} bond);

R^{10} is H, CH_3 , CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH_2O-R^{28} or $-OR^{28}$, where R^{28} is H or together with R^4 forms a bond (lactone) or R^{10} together with R^1 forms a bond (C_1-C_{10} double bond);

R^{11} is H, or OH or is absent;

R^{12} is CH_3 , CH_2OH , $COOH$ or a glycosylic ester (glycoside ester) of said $COOH$;

R^{13} is methylene, or a divalent hetero-atom, or NR^{29} , where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl; and a double bond is present between C_{16} and R^{13} when R^{11} is absent; or R^{13} is H, OH, CH_3 CHO, CH_2X , where X is halogen, $CHNR^{29}$ where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl when R^{11} is H or OH; with the proviso that where R^{11} is OH, R^{13} is not OH;

R^{14} is H or OH;

R^{15} is H, or together with R^9 forms a bond (C_9 - C_{15} bond);


and its pharmaceutically acceptable **lactones, esters, active esters, salts and organic bases, derivatives**

in combination with other compatible therapeutic agents selected from the group consisting of analgesics, anti-hypertensive agents, sedatives, hypnotics, lipid-lowering agents, and anti-infective agents or combinations thereof, to a patient in need thereof.

9. (Currently Amended) A method according to claim **11** **[[8]]**, wherein the Gibberellins are Gibberellin A_3 .

10. (Currently Amended) A method according to claim **11** **[[8]]**, wherein the Gibberellins are a mixture of Gibberellin A_3 and Gibberellin A_4 and/or Gibberellin A_7 .

11. (Currently amended) A method of treatment **for Type I and Type II** of diabetes and **its complications and associated** related conditions comprising administering compounds **selected from** of formula (1) (Gibberellins)



Formula (1)

wherein

R^1 is H or a group $-O-R^{20}$, where R^{20} is H, a glycosylic ether group (glycoside ether), C_{1-6} alkyl group, or R^1 together with R^2 or R^{10} forms a bond (C_1-C_2 or C_1-C_{10} double bond, respectively);

R^2 is H or a group $-O-R^{21}$, where R^{21} is H, a glycosylic ether group (glycoside ether), or together with R^4 forms a bond (lactone) or R^2 together with R^1 or R^3 forms a bond (C_1-C_2 or C_2-C_3 double bond, respectively);

R^3 is H, $=O$, or $-O-R^{22}$, where R^{22} is H or a glycosylic ether group (glycoside ether), or R^3 together with R^2 forms a bond (C_2-C_3 double bond);

R^4 is OH, or $-OR^{23}$, where R^{23} is unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, arylalkyl, amidine, $-NR^{24}R^{25}$ or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur; R^{24} and R^{25} may or may not be the same, are hydrogen, or C_{1-20} alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulphur; or R^4 together with R^{21} or R^{28} forms a bond (lactone);

R⁵ is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted C₁₋₂₀ alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters;

R⁶ is H or OH or together with R⁷ forms a bond (C₁₁-C₁₂ double bond);

R⁷ is H, =O, or -OR²⁶, where R²⁶ is H or a glycosylic ether group (glycoside ether) or R⁷ together with R⁶ forms a bond (C₁₁-C₁₂ double bond);

R⁸ is H, hydroxyl, mercaptan, or halogen, amino, azido, NR²⁴R²⁵, unsubstituted or substituted C₁₋₂₀ alkyl, allyl, aryl, or arylalkyl, or -OR²⁷, where R²⁷ is a glycosylic ether group (glycoside ether);

R⁹ is H or OH, or together with R¹⁵ forms a bond (C₉-C₁₅ bond);

R¹⁰ is H, CH₃, CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH₂O-R²⁸ or -OR²⁸, where R²⁸ is H or together with R⁴ forms a bond (lactone) or R¹⁰ together with R¹ forms a bond (C₁-C₁₀ double bond);

R¹¹ is H, or OH or is absent;

R¹² is CH₃, CH₂OH, COOH or a glycosylic ester (glycoside ester) of said COOH;

R¹³ is methylene, or a divalent hetero-atom, or NR²⁹, where R²⁹ is NHR³⁰ or OR³⁰ where R³⁰ is H, or C₁₋₂₀ alkyl, aryl, alkylaryl; and a double bond is present between C₁₆ and R¹³ when R¹¹ is absent; or R¹³ is H, OH, CH₃, CHO, CH₂X, where X is halogen, CHNR²⁹ where R²⁹ is NHR³⁰ or OR³⁰ where R³⁰ is H, or C₁₋₂₀ alkyl, aryl, alkylaryl when R¹¹ is H or OH; with the proviso that where R¹¹ is OH, R¹³ is not OH;


R¹⁴ is H or OH;

R^{15} is H, or together with R^9 forms a bond (C_9 - C_{15} bond);

and or their pharmaceutically acceptable lactones, esters, active esters, salts and organic bases derivatives,

in combination with substances selected from the group consisting of insulin, its fragment derivatives, IGFs, growth factors, and other pharmaceutically compatible anti-diabetic agents, or combinations thereof, to a patient in need thereof.

12. (Currently amended) A method of treatment for Type I and Type II of diabetes and its complications and associated related conditions comprising administering compounds selected from of formula (1) (Gibberellins)



Formula (1)

wherein

R^1 is H or a group $-O-R^{20}$, where R^{20} is H, a glycosylic ether group (glycoside ether), C_{1-6} alkyl group, or R^1 together with R^2 or R^{10} forms a bond (C_1 - C_2 or C_1 - C_{10} double bond, respectively);

R^2 is H or a group $-O-R^{21}$, where R^{21} is H, a glycosylic ether group (glycoside ether), or together with R^4 forms a bond (lactone) or R^2 together with R^1 or R^3 forms a bond (C_1-C_2 or C_2-C_3 double bond, respectively);

R^3 is H, $=O$, or $-O-R^{22}$, where R^{22} is H or a glycosylic ether group (glycoside ether), or R^3 together with R^2 forms a bond (C_2-C_3 double bond);

R^4 is OH, or $-OR^{23}$, where R^{23} is unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, arylalkyl, amidine, $-NR^{24}R^{25}$ or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur; R^{24} and R^{25} may or may not be the same, are hydrogen, or C_{1-20} alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulphur; or R^4 together with R^{21} or R^{28} forms a bond (lactone);

R^5 is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted C_{1-20} alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters;

R^6 is H or OH or together with R^7 forms a bond ($C_{11}-C_{12}$ double bond);

R^7 is H, $=O$, or $-OR^{26}$, where R^{26} is H or a glycosylic ether group (glycoside ether) or R^7 together with R^6 forms a bond ($C_{11}-C_{12}$ double bond);

R^8 is H, hydroxyl, mercaptan, or halogen, amino, azido, $NR^{24}R^{25}$, unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, or arylalkyl, or $-OR^{27}$, where R^{27} is a glycosylic ether group (glycoside ether);

R^9 is H or OH, or together with R^{15} forms a bond (C_9-C_{15} bond);

R^{10} is H, CH_3 , CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH_2O-R^{28} or $-OR^{28}$, where R^{28} is H or together with R^4 forms a bond (lactone) or R^{10} together with R^1 forms a bond (C_1 - C_{10} double bond);

R^{11} is H, or OH or is absent;

R^{12} is CH_3 , CH_2OH , COOH or a glycosylic ester (glycoside ester) of said COOH;

R^{13} is methylene, or a divalent hetero-atom, or NR^{29} , where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl; and a double bond is present between C_{16} and R^{13} when R^{11} is absent; or R^{13} is H, OH, CH_3 , CHO, CH_2X , where X is halogen, $CHNR^{29}$ where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl when R^{11} is H or OH; with the proviso that where R^{11} is OH, R^{13} is not OH;

R^{14} is H or OH;

R^{15} is H, or together with R^9 forms a bond (C_9 - C_{15} bond);

and its or their pharmaceutically acceptable lactones, esters, active esters, salts and organic bases derivatives,

in combination with **substances selected from the group consisting of insulin, its fragment derivatives, IGFs, growth factors, and other pharmaceutically compatible anti-diabetic agents, or combinations thereof, along with** other compatible therapeutic agents selected from the group consisting of analgesics, anti-hypertensive agents, sedatives, hypnotics, lipid-lowering agents, and anti-infective agents or combinations thereof, to a patient in need thereof.


13. (Currently amended) The method according to claim **11** **[[1]]**, for the treatment of type 1 diabetes and its associated conditions.

14. (Currently amended) The method according to claim 11 [[1]], for the treatment of type 2 diabetes and its associated conditions.

15. (Currently amended) The method according to claim 14 [[1]], for the treatment of insulin resistant diabetes.

16. (Currently amended) The method according to claim 1, wherein the diabetic related complications and associated conditions [[,]] are chosen from obesity, micro and macro vascular diseases, nephropathy, neuropathy and eye diseases.

17. (Currently amended) An anti-diabetic agent consisting essentially of comprising a compound of formula (1)



Formula (1)

wherein

R¹ is H or a group -O-R²⁰, where R²⁰ is H, a glycosylic ether group (glycoside ether), C₁₋₆ alkyl group, or R¹ together with R² or R¹⁰ forms a bond (C₁-C₂ or C₁-C₁₀ double bond, respectively);

R^2 is H or a group $-O-R^{21}$, where R^{21} is H, a glycosylic ether group (glycoside ether), or together with R^4 forms a bond (lactone) or R^2 together with R^1 or R^3 forms a bond (C_1-C_2 or C_2-C_3 double bond, respectively);

R^3 is H, $=O$, or $-O-R^{22}$, where R^{22} is H or a glycosylic ether group (glycoside ether), or R^3 together with R^2 forms a bond (C_2-C_3 double bond);

R^4 is OH, or $-OR^{23}$, where R^{23} is unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, arylalkyl, amidine, $-NR^{24}R^{25}$ or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur; R^{24} and R^{25} may or may not be the same, are hydrogen, or C_{1-20} alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulphur; or R^4 together with R^{21} or R^{28} forms a bond (lactone);

R^5 is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted C_{1-20} alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters;

R^6 is H or OH or together with R^7 forms a bond ($C_{11}-C_{12}$ double bond);

R^7 is H, $=O$, or $-OR^{26}$, where R^{26} is H or a glycosylic ether group (glycoside ether) or R^7 together with R^6 forms a bond ($C_{11}-C_{12}$ double bond);

R^8 is H, hydroxyl, mercaptan, or halogen, amino, azido, $NR^{24}R^{25}$, unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, or arylalkyl, or $-OR^{27}$, where R^{27} is a glycosylic ether group (glycoside ether);

R^9 is H or OH, or together with R^{15} forms a bond (C_9-C_{15} bond);

R^{10} is H, CH_3 , CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH_2O-R^{28} or $-OR^{28}$, where R^{28} is H or together with R^4 forms a bond (lactone) or R^{10} together with R^1 forms a bond (C_1-C_{10} double bond);

R^{11} is H, or OH or is absent;

R^{12} is CH_3 , CH_2OH , COOH or a glycosylic ester (glycoside ester) of said COOH;

R^{13} is methylene, or a divalent hetero-atom, or NR^{29} , where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl; and a double bond is present between C_{16} and R^{13} when R^{11} is absent; or R^{13} is H, OH, CH_3 , CHO, CH_2X , where X is halogen, $CHNR^{29}$ where R^{29} is NHR^{30} or OR^{30} where R^{30} is H, or C_{1-20} alkyl, aryl, alkylaryl when R^{11} is H or OH; with the proviso that where R^{11} is OH, R^{13} is not OH;


R^{14} is H or OH;

R^{15} is H, or together with R^9 forms a bond (C_9-C_{15} bond);

and/or its **pharmaceutically acceptable lactones, esters, active esters, salts and organic bases** derivatives as an active ingredient, together with a pharmaceutically acceptable carrier.

18. (Original) An anti-diabetic agent according to claim 17, wherein the agent is a medicament suitable for administration with a medicator.

19. (Currently amended) An anti-diabetic agent **consisting essentially of** comprising a compound of formula (1)



Formula (1)

wherein

R^1 is H or a group $-O-R^{20}$, where R^{20} is H, a glycosylic ether group (glycoside ether), C_{1-6} alkyl group, or R^1 together with R^2 or R^{10} forms a bond (C_1-C_2 or C_1-C_{10} double bond, respectively);

R^2 is H or a group $-O-R^{21}$, where R^{21} is H, a glycosylic ether group (glycoside ether), or together with R^4 forms a bond (lactone) or R^2 together with R^1 or R^3 forms a bond (C_1-C_2 or C_2-C_3 double bond, respectively);

R^3 is H, $=O$, or $-O-R^{22}$, where R^{22} is H or a glycosylic ether group (glycoside ether), or R^3 together with R^2 forms a bond (C_2-C_3 double bond);

R^4 is OH, or $-OR^{23}$, where R^{23} is unsubstituted or substituted C_{1-20} alkyl, allyl, aryl, arylalkyl, amidine, $-NR^{24}R^{25}$ or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur; R^{24} and R^{25} may or may not be the same, are hydrogen, or C_{1-20} alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulphur; or R^4 together with R^{21} or R^{28} forms a bond (lactone);

R⁵ is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted C₁₋₂₀ alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters;

R⁶ is H or OH or together with R⁷ forms a bond (C₁₁-C₁₂ double bond);

R⁷ is H, =O, or -OR²⁶, where R²⁶ is H or a glycosylic ether group (glycoside ether) or R⁷ together with R⁶ forms a bond (C₁₁-C₁₂ double bond);

R⁸ is H, hydroxyl, mercaptan, or halogen, amino, azido, NR²⁴R²⁵, unsubstituted or substituted C₁₋₂₀ alkyl, allyl, aryl, or arylalkyl, or -OR²⁷, where R²⁷ is a glycosylic ether group (glycoside ether);

R⁹ is H or OH, or together with R¹⁵ forms a bond (C₉-C₁₅ bond);

R¹⁰ is H, CH₃, CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH₂O-R²⁸ or -OR²⁸, where R²⁸ is H or together with R⁴ forms a bond (lactone) or R¹⁰ together with R¹ forms a bond (C₁-C₁₀ double bond);

R¹¹ is H, or OH or is absent;

R¹² is CH₃, CH₂OH, COOH or a glycosylic ester (glycoside ester) of said COOH;

R¹³ is methylene, or a divalent hetero-atom, or NR²⁹, where R²⁹ is NHR³⁰ or OR³⁰ where R³⁰ is H, or C₁₋₂₀ alkyl, aryl, alkylaryl; and a double bond is present between C₁₆ and R¹³ when R¹¹ is absent; or R¹³ is H, OH, CH₃, CHO, CH₂X, where X is halogen, CHNR²⁹ where R²⁹ is NHR³⁰ or OR³⁰ where R³⁰ is H, or C₁₋₂₀ alkyl, aryl, alkylaryl when R¹¹ is H or OH; with the proviso that where R¹¹ is OH, R¹³ is not OH;

R¹⁴ is H or OH;

R¹⁵ is H, or together with R⁹ forms a bond (C₉-C₁₅ bond);

and/or its pharmaceutically acceptable lactones, esters, active esters, salts and organic bases as an active ingredient, together with pharmaceutically acceptable carriers or excipients, wherein the agent is a slow release composition.

20. (Original) An anti-diabetic agent according to claim 17, wherein the agent is for oral administration.

21. (Original) An anti-diabetic agent according to claim 17, wherein the agent is for inhalation administration.

22. (Original) An anti-diabetic agent according to claim 17, wherein the agent is for transdermal administration.

23. (Original) An anti-diabetic agent according to claim 17, wherein the agent is for parenteral injection.

24. (Original) An anti-diabetic agent according to claim 17, wherein the agent is for topical, rectal, or vaginal administration.

25. (Canceled)

26. (Currently amended) An anti-diabetic agent according to claim 17, ~~[[25]]~~ wherein the pharmaceutically acceptable salt derivative is a sodium salt of formula (1).

27. (Currently amended) An anti-diabetic agent according to claim 17, ~~[[25]]~~ wherein the pharmaceutically acceptable salt derivative is a zinc salt of formula (1).

28. (Currently amended) An anti-diabetic agent according to claim 17, ~~[[25]]~~ wherein the pharmaceutically acceptable ester derivative is a ethyl ester of formula (1).

29. (Currently amended) A method of manufacturing an anti-diabetic agent according to claim 17, comprising combining a compound selected from of formula (1) and ~~and/or~~ its pharmaceutically acceptable lactones, esters, active esters, salts and organic bases derivatives with a pharmaceutically acceptable carrier.

30. (Withdrawn) A process for the preparation of Gibberellins including Gibberellin A₃, including the steps of:

- (a) incubating a Gibberellin-producing strain of microorganism in a fermentation broth;
- (b) adjusting the pH of the fermentation broth to pH 6.5 to 7.0 and filtering to obtain a filter cake of microorganism mycelium, and a filtrate;
- (c) washing the filter cake with water and combining the washing with the filtrate to form an aqueous solution;
- (d) concentrating the aqueous solution;
- (e) mixing the aqueous solution with an organic solvent at a temperature of 5 to 10°C and adjusting the pH of the mixture to less than 2.0;
- (f) allowing the mixture to separate into an aqueous phase and a first organic phase and removing the first organic phase;
- (g) re-extracting the aqueous phase from step (f) with organic solvent to obtain a second organic phase;
- (h) combining the first and second organic phases and concentrating to form a concentrated organic solution;
- (i) heating the concentrated organic solution at 60-70°C for 3 to 4 hours with stirring, until the precipitation of solid matter ceases;

- (j) cooling the concentrated organic solution to room temperature and filtering to obtain a precipitate;
- (k) washing the precipitate in cold organic solvent and drying to obtain an off-white solid containing about 80% Gibberellin A₃, about 4% Gibberellin A₄ and about 4% Gibberellin A₇.

31. (Withdrawn) The process of claim 30, comprising the further steps of:

- (l) dissolving the off-white solid in a mixture of 32.6% methanol, 2.2% water and 65.2% acetone to obtain a Gibberellin solution;
- (m) diluting the Gibberellin solution with a 10:1 mixture of organic solvent and water;
- (n) filtering the diluted Gibberellin solution and concentrating the filtrate by vacuum evaporation;
- (o) heating the concentrate to a temperature of 60 to 80°C for 2 to 3 hours with stirring, cooling to room temperature and filtering to obtain a solid crystalline precipitate;
- (p) washing the precipitate with cold organic solvent and drying to obtain Gibberellin A₃ crystals.

32. (Withdrawn) A process according to claim 30 wherein the Gibberellin-producing strain of microorganism is *Gibberella fujikuroi*.

33. (Withdrawn) A process according to claim 30, wherein the concentration of the solutions in steps (d) and (h) is achieved using vacuum evaporation.

34. (Withdrawn) A process according to claim 30 wherein the organic solvent is ethyl acetate.

35. (Withdrawn) A process according to claim 31 wherein the organic solvent is ethyl acetate.

36. (Withdrawn) A process according to claim 31 comprising the further steps of:

- (q) dissolving the Gibberellin A₃ in methanol;
- (r) adding the Gibberellin solution to an equimolar aqueous solution of NaHCO₃;
- (s) evaporating the mixed solutions to dryness to obtain a solid residue;
- (t) dissolving the residue in water and freeze drying to obtain Gibberellin A₃ sodium salt.

37. (Withdrawn) A process according to claim 36, comprising the further steps of dissolving the Gibberellin A₃ sodium salt in water, passing the solution through a column loaded with a zinc ion-exchange resin, washing the column with water, collecting and combining the effluent and washings and removing the water to obtain Gibberellin A₃ zinc salt.

38. (Withdrawn) A process according to claim 31 comprising the further steps of:

- (q) dissolving the Gibberellin A₃ in a 50:1 ratio mixture of acetone to water;
- (r) mixing the Gibberellin A₃ solution with equimolar amounts of triethylamine and ethyl chloroformate, and a one tenth molar amount of N-methyl morpholine, and stirring at -15°C for 20 minutes;
- (s) diluting the resultant mixture with anhydrous ethanol and stirring at room temperature;
- (t) evaporating the diluted mixture to dryness and partitioning the residue between ethyl acetate and water in a 6:1 ratio;
separating the ethyl acetate layer, washing with 2% HCl, followed by water, followed by 5% NaHCO₃, followed by water, and evaporating under reduced pressure to dryness to give Gibberellin A₃ ethyl ester.

39. (New) The method of claim 11, wherein the complications and associated conditions of diabetes are one or more of: obesity, micro and macro vascular diseases, nephropathy, neuropathy, eye diseases, and diabetic ulcerations.

40. (New) The method of claim 11, wherein the pharmaceutically acceptable salts are selected from alkali metal salts, alkaline earth metal salts, metal, and salts of ammonium or organic bases.

41. (New) The method of claim 40, wherein the organic bases are lidocaine, or $\text{NR}^{16} \text{R}^{17} \text{R}^{18} \text{R}^{19}$, where R^{16} , R^{17} , R^{18} , R^{19} , which may be the same or not the same, are hydrogen, or substituted or unsubstituted C_{1-20} alkyl, alkanol, or aryl groups.